# Corpus Christi Institute of Cosmetic & Plastic Surgery

## POST OPERATIVE INSTRUCTIONS FOR CAPSULECTOMY

### **GENERAL INSTRUCTIONS**

- Have someone drive you home after surgery and help you at home for 1-2 days.
- Get plenty of rest and increase your fluid intake.
- Decreased activity may promote constipation, so you may want to add more ripe fruit to your diet. Prunes and prune juice work well for most.
- Take pain medications as prescribed. Do not take aspirin or any products containing aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs).
- Do not drink alcohol when taking pain medications. •
- Refrain from drinking alcohol for three weeks following surgery, even when no longer • taking pain medications, as alcohol causes fluid retention.
- If you regularly take vitamins with iron, resume these as tolerated.
- Do not smoke or consume nicotine, as smoking delays healing and increases the risk of • complications.
- Please get Arnica in the pill form. Take once a day starting the day of surgery once you get home. Continue taking it for 2 weeks post op. This will help with bruising.

## **ACTIVITY RESTRICTIONS**

- Start walking as soon as possible with assistance the day of surgery, this helps to reduce • swelling and lowers the chance of blood clots. Starting the day after surgery, walk 300 feet three times per day with assistance daily.
- Do not drive until you are no longer taking any pain medications (narcotics). •
- Do not drive until you have full range of motion with your arms. •
- Refrain from vigorous activity for two weeks. •
- Body contact sports should be avoided for 6-8 weeks.
- Social and employment activities can be resumed in 3-10 days. •
- Sexual activity can be resumed in two weeks unless otherwise specified by your • surgeon.

## **INCISION CARE**

- You may shower 48 hours after surgery.
- Keep steri-strips on, and replace if they come off. Keep incisions clean and inspect daily for sign of infection. No soaking incision with water for two weeks after surgery.
- You may pad the incisions with gauze for comfort.

Corpus Christi Institute of Cosmetic & Plastic Surgery Initials \_\_\_\_\_ 5642 Esplanade Drive Corpus Christi, Texas 78414 Phone: (361) 888-7417 - Fax: (361) 651-1489

# Corpus Christi Institute of Cosmetic & Plastic Surgery

## POST OPERATIVE INSTRUCTIONS FOR CAPSULECTOMY

- Wear garments (bra, bandeau) as directed by surgeon. Do not wear underwire bra for six weeks after surgery.
- Refrain from sleeping on your stomach for three weeks. •

## **EXPECTATIONS**

- Expect some drainage onto the steri-strips covering the incisions.
- You are likely to feel tired for a few days. •
- Maximum discomfort will occur in the first few days after surgery. •
- You may experience some numbness of nipples and operative areas.
- You may experience a burning sensation in your nipples for about two weeks.
- You may experience temporary soreness tightness, swelling, and bruising as well as • some discomfort in the incision area.
- Your breasts may be sensitive to stimulation for a few weeks. •
- You may feel enlarged as if you are ready to breast feed. This should resolve in a few weeks.

## APPEARANCE AND SCAR CARE

- Most of the discoloration and swelling will subside in 4-6 weeks. •
- Scars may be red and angry looking for six months. In time, these usually soften and • fade.
- Avoid exposure of scar to sunlight for 6-12 months. If unavoidable, use sunblock (SPF 30 • or greater) with zinc oxide or titanium dioxide.
- Massage scars ten minutes at a time, three times daily while using silicone strips. ٠

## FOLLOW UP CARE

Your post-operative appointment will be scheduled before surgery for your • convenience.

#### WHEN TO CALL

- If you have increased swelling or bruising.
- If swelling and redness persist after a few days.
- If you have increased redness along the incision.
- If you have severe or increased pain not relieved by medication. •
- If you have any side effects such as rash, nausea, headache, or vomiting. •
- If you have an oral temperature over 100.4 degrees. •
- If you have yellowish or greenish drainage from the incisions or notice a foul odor. •
- If you have bleeding from the incisions that is difficult to control with light pressure. •
- If you have loss of feeling or motion. •

Corpus Christi Institute of Cosmetic & Plastic Surgery Initials 5642 Esplanade Drive Corpus Christi, Texas 78414 Phone: (361) 888-7417 - Fax: (361) 651-1489