

Corpus Christi Institute of Cosmetic & Plastic Surgery

POST OPERATIVE INSTRUCTIONS FOR LIPOSUCTION

GENERAL INSTRUCTIONS

- Have someone drive you home after surgery and help you at home for 1-2 days.
- Get plenty of rest and increase your fluid intake.
- Decreased activity may promote constipation, so you may want to add more ripe fruit to your diet. Prunes and prune juice work well for most.
- Take pain medications as prescribed. Do not take aspirin or any products containing aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs).
- Do not drink alcohol when taking pain medications.
- Refrain from drinking alcohol for three weeks following surgery, even when no longer taking pain medications, as alcohol causes fluid retention.
- If you regularly take vitamins with iron, resume these as tolerated.
- Do not smoke or consume nicotine, as smoking delays healing and increases the risk of complications.
- **Please get Arnica in the pill form. Take once a day starting the day of surgery once you get home. Continue taking it for 2 weeks post op. This will help with bruising.**

ACTIVITY RESTRICTIONS

- Start walking as soon as possible with assistance the day of surgery, this helps to reduce swelling and lowers the chance of blood clots. Starting the day after surgery, walk 300 feet three times per day with assistance daily.
- Do not drive until you are no longer taking any pain medications (narcotics).
- Activities involving affected sites should be minimized for approximately 2 weeks. Strenuous exercise and activities should be avoided for 4-6 weeks.
- Sexual activity can be resumed in two weeks unless otherwise specified by your surgeon.

INCISION CARE

- You may shower 48 hours after surgery.
- Keep steri-strips on, and replace if they come off. Keep incisions clean and inspect daily for sign of infection. No tub soaking while sutures or drains are in place.
- Place daily soft dressing over incisions and around drain sites to wick away moisture and prevent irritation by garment along the incision line.
- Wear compression garment all the time (24/7) for six weeks post operatively, unless showering.

Initials_____

Corpus Christi Institute of Cosmetic & Plastic Surgery
5642 Esplanade Drive
Corpus Christi, Texas 78414
Phone: (361) 888-7417 - Fax: (361) 651-1489

Corpus Christi Institute of Cosmetic & Plastic Surgery

POST OPERATIVE INSTRUCTIONS FOR LIPOSUCTION

EXPECTATIONS

- You will have small incision sites at the point of introduction of the instrument used to remove the fat tissue.
- The small incision sites will be covered with gauze to protect these areas.
- There will be swelling and discoloration after surgery.
- There may be numbness, bruising, swelling, soreness, burning sensation, bleeding, pain and discomfort around the operative areas.
- Most of the swelling and discoloration usually subsides in 6-8 weeks, but some may persist for 6 months or more.
- May have fluid drainage from incision sites.
- A drainage tube may be inserted beneath the skin to prevent fluid build-up.
- It is normal to feel a bit anxious or depressed in the days or weeks following surgery; but, generally, after about a week, you should begin to feel better.
- Bruising may be apparent beyond the areas of liposuction, partially due to gravity.

APPEARANCE AND DRAIN CARE

- Improved skin texture, firmer, and smoother skin.
- Empty drain bulb 3 times a day and record outputs in mL (milliliters) noting time of day and drain amount. Make sure to record each drain's output separately.
- It is normal for drain outputs to vary and for clots to be in tube and bulb.
- **Do not disconnect bulb from tube.**
- **Do not push in or pull on the drain tube at its entrance on the skin.**
- See additional attachment for more instructions and additional drain log.

FOLLOW UP CARE

- Your post-operative appointment will be scheduled before surgery for your convenience.

WHEN TO CALL

- If you have increased swelling or bruising.
- If swelling and redness persist after a few days.
- If you have increased redness along the incision.
- If you have severe or increased pain not relieved by medication.
- If you have any side effects such as rash, nausea, headache, or vomiting.
- If you have an oral temperature over 100.4 degrees.
- If you have yellowish or greenish drainage from the incisions or notice a foul odor.
- If you have bleeding from the incisions that is difficult to control with light pressure.
- If you have loss of feeling or motion.

Initials_____

Corpus Christi Institute of Cosmetic & Plastic Surgery
5642 Esplanade Drive
Corpus Christi, Texas 78414
Phone: (361) 888-7417 - Fax: (361) 651-1489